

(L1) in feces



9/1/


Parascaris equorum	
Common Name	Equine roundworm
Infective Stage	Second-stage larva (L2)
Habitat	Small intestine
Mode of transmission	Ingesting embryonated eggs
	containing L2 and L2 emerge
	from the eggs
Diagnostic Specimen	Feces
Diagnostic State	Egg in feces / Adult worms in
10	the intestine



Strongylus edentatus	
Common Name	Equine roundworm
Infective Stage	Third-stage larva (L3)
Habitat	Large intestine (caecum / colon)
Mode of transmission	Ingestion of Larva 3
Diagnostic Specimen	Feces / subperitoneum
Diagnostic State	Egg in feces or immature adult
0 00	and L4 cause subperitoneal cysts
U V	ที่บริเวณสวาป



Strongylus equines	
Common Name	Equine roundworm
Infective Stage	Third-stage larva (L3)
Habitat	Large intestine (caecum / colon)
Mode of transmission	Ingestion of Larva 3
Diagnostic Specimen	Feces / subperitoneum
Diagnostic State	Egg in feces or immature adult
9 6	and L4 cause subperitoneal cysts
10	ที่บริเวณสวาป



Strongylus vulgaris	
Common Name	Equine roundworm
Infective Stage	Third-stage larva (L3)
Habitat	Large intestine (caecum / colon)
Mode of transmission	Ingestion of Larva 3
Diagnostic Specimen	Feces / subperitoneum
Diagnostic State	Egg in feces

